

Quantum Langlands duality and mirror symmetry

Igor Yu. Potemine

1 Introduction

There are recent indications that geometric Langlands (Beilinson-Drinfeld) duality [2, 3] is related to S -duality in QFT and string theories (*cf.* [30]). In this paper we discuss a peculiar relation of this Beilinson-Drinfeld duality and Givental's version of mirror symmetry. It should be noted that mirror symmetry is a T -duality from the physical viewpoint [29].

Consider two (infinite-dimensional) ind-varieties: the n -*reduced Sato Grassmannian* $\mathcal{GR}^{(n)}$ and the *periodic* (or *affine*) *flag variety* $\mathcal{FL}^{(n)}$. The *Sato Grassmannian* $\mathcal{GR}$ is the set of subspaces $W \subset \mathbb{C}((t))$ commensurable with $\mathbb{C}[t^{-1}]$, *i. e.*, such that the projection

$$\gamma_W : W \rightarrow \mathbb{C}((t))/t\mathbb{C}[[t]] \quad (1.1)$$

is Fredholm (with finite kernel and cokernel). Then

$$\mathcal{GR}^{(n)} = \{W \in \mathcal{GR} \mid t^{-n}W \subset W\} \quad (1.2)$$

and

$$\mathcal{FL}^{(n)} = \{W_0 \subset W_1 \subset \dots \subset W_n \mid \forall i \ W_i \in \mathcal{GR}^{(n)}, \\ \text{virt. dim. } W_i = i - n \text{ and } t^{-n}W_n = W_0\} \quad (1.3)$$

where

$$\text{virt. dim. } W = \text{ind } \gamma_W = \dim \text{Ker } \gamma_W - \dim \text{Coker } \gamma_W \quad (1.4)$$

is the *virtual dimension* of W .

We construct $\mathcal{D}$ -modules of Beilinson-Drinfeld type on $\mathcal{GR}^{(n)}$ parametrized by local affine GL_n -opers. In the same manner, we construct $\mathcal{D}$ -modules of Givental type on $\mathcal{FL}^{(n)}$ parametrized by local affine Miura GL_n -opers.

MAIN THEOREM. *Givental (quantum) $\mathcal{D}$ -modules on $\mathcal{FL}^{(n)}$ are transformed to Beilinson-Drinfeld (quantum) $\mathcal{D}$ -modules on $\mathcal{GR}^{(n)}$ via the affine Miura transformation.*

The outline of this paper is as follows. First of all, we recall some facts about the gravitational quantum cohomology and Givental's construction of a quantum $\mathcal{D}$ -module $\mathcal{D}(X)$ for a smooth projective variety X . We would like to construct a family of $\mathcal{D}$ -modules on infinite-dimensional varieties $\mathfrak{GR}^{(n)}$ and $\mathcal{FL}^{(n)}$ parametrized by affineopers. It is possible to generalize Givental's approach using Guest-Otofugi results [14]. However, in this paper we adopt a different strategy. In section 4 we recall Beilinson-Drinfeld construction of $\mathcal{D}$ -modules on the moduli stack $\text{Bun}_G(X)$ of principal G -bundles on a curve X . This construction uses a quantization of the Hitchin system. In the following two sections (5 and 6) we sketch a local affine version of the Beilinson-Drinfeld construction using the formalism of null vectors. In some sense, it is related to the quantization of Korteweg-de Vries hierarchies. A similar approach is used later to construct $\mathcal{D}$ -modules on $\mathcal{FL}^{(n)}$ parametrized by local affine Miura GL_n -opers. In section 9 we deduce our main theorem from standard results about the Miura transformation. Finally, we indicate how one can follow very closely the Beilinson-Drinfeld construction using analogues of the Hitchin fibration.

2 Background from gravitational quantum cohomology

The surveys [5] and [23] may serve as a nice introduction.

Physically, (gravitational) quantum cohomology is a cohomological realization of non-linear Σ -models of two-dimensional field theory (coupled to gravity). One considers interacting “strings” propagating inside a variety X , *i. e.*, maps $\Sigma \rightarrow X$ where Σ is a Riemann surface swept out by these strings. The main statistical objects of interest are the *correlators* (vacuum expectation values) of the *primary fields*, represented by cocycles $\gamma_1, \dots, \gamma_n \in H^*(X, \mathbb{Q})$, and their *gravitational descendants* $\tau_{d_1}\gamma_1, \dots, \tau_{d_n}\gamma_n$ ($d_i \in \mathbb{N}$). Fix a homology class $\beta \in H_2(X, \mathbb{Z})$ and consider the moduli stack $\overline{\mathcal{M}}_{g,n}(X, \beta)$ of stable maps $f : C \rightarrow X$ of algebraic curves of genus g with n marked points such that $f_*[C] = \beta$. Denote by $[\overline{\mathcal{M}}_{g,n}(X, \beta)]^{\text{virt}}$ the *virtual fundamental class* (the orbifoldic Poincaré dual of $1 \in H^*(\overline{\mathcal{M}}_{g,n}(X, \beta), \mathbb{Q})$). In addition, let

$$e_i : \overline{\mathcal{M}}_{g,n}(X, \beta) \longrightarrow X, \quad e_i(f : C \rightarrow X, p_1, \dots, p_n) = f(p_i) \quad (2.1)$$

be the evaluation maps at i th marked point and $\mathcal{L}_i$ “the cotangent line at p_i ”, *i. e.*, a line bundle over $\overline{\mathcal{M}}_{g,n}(X, \beta)$ whose fiber over $(f : C \rightarrow X, p_1, \dots, p_n)$ is the cotangent space $T_{p_i}^*C$. Now the *gravitational correlators* are defined by

$$\langle \tau_{d_1}\gamma_1, \dots, \tau_{d_n}\gamma_n \rangle_{g,\beta} = \int_{\overline{\mathcal{M}}_{g,n}(X, \beta)} I_{g,n,\beta}(\tau_{d_1}\gamma_1, \dots, \tau_{d_n}\gamma_n)$$

where

$$I_{g,n,\beta}(\tau_{d_1}\gamma_1, \dots, \tau_{d_n}\gamma_n) = (c_1(\mathcal{L}_1)^{d_1} \cup e_1^*(\gamma_1)) \cup \dots \cup (c_1(\mathcal{L}_n)^{d_n} \cup e_n^*(\gamma_n))$$

are the *gravitational Gromov-Witten classes*. If $d_1 = \dots = d_n = 0$ then

$$I_{g,n,\beta}(\tau_{d_1}\gamma_1, \dots, \tau_{d_n}\gamma_n) = I_{g,n,\beta}(\gamma_1, \dots, \gamma_n) \quad (2.2)$$

are the usual *Gromov-Witten classes* and

$$\langle \tau_{d_1}\gamma_1, \dots, \tau_{d_n}\gamma_n \rangle_{g,\beta} = \langle I_{g,n,\beta} \rangle(\gamma_1, \dots, \gamma_n) \quad (2.3)$$

are the *Gromov-Witten invariants*.

3 Givental quantum $\mathcal{D}$ -modules

First of all, we recall some basic facts about Givental quantum connection and associated quantum $\mathcal{D}$ -modules. Let X be a smooth projective variety and $T_0, \dots, T_m$ a basis of $H^*(X, \mathbb{C})$. We consider $H^*(X, \mathbb{C})$ as a Frobenius supermanifold with respect to supercommutative variables $t_0, \dots, t_m$ associated to T_i (cf. [5, 8.2.1]). The *Givental quantum connection* ∇^h is defined on the trivial cohomology bundle $H^*(X, \mathbb{C}[[t_0, \dots, t_m]])$ over $H^*(X, \mathbb{C})$ by

$$\nabla_{\frac{\partial}{\partial t_i}}^h \left(\sum_j a_j T_j \right) = \hbar \sum_j \frac{\partial a_j}{\partial t_i} T_j - \sum_j a_j T_j * T_i \quad (3.1)$$

where $*$ is the *big quantum product* on $H^*(X, \mathbb{C}[[t_0, \dots, t_m]])$.

Consider the *genus g couplings*

$$\langle \langle \tau_{d_1}\gamma_1, \dots, \tau_{d_n}\gamma_n \rangle \rangle_g = \sum_{k=0}^{\infty} \sum_{\beta} \frac{1}{k!} \langle \tau_{d_1}\gamma_1, \dots, \tau_{d_n}\gamma_n, \underbrace{\gamma, \dots, \gamma}_{k \text{ times}} \rangle_{g,\beta} q^\beta \quad (3.2)$$

where $\gamma = \sum_i t_i T_i$ [5, sect. 10.1.1]. The formal sections

$$s_a = T_a + \sum_j \left\langle \left\langle \frac{T_a}{\hbar - c}, T_j \right\rangle \right\rangle_0 T^j, \quad (3.3)$$

where $c = c_1(\mathcal{L}_1)$, form a basis of ∇^h -flat sections of $H^*(X, \mathbb{C}[[t_0, \dots, t_m]])$. This means that

$$\hbar \frac{\partial s_a}{\partial t_i} = T_i * s_a, \quad a, i = 0, \dots, m \quad (3.4)$$

[5, sect. 10.2.1].

DEFINITION 3.1 Denote

$$\mathcal{D} = \mathbb{C}[\hbar \partial / \partial t_i, \exp t_i, \hbar]. \quad (3.5)$$

The elements of $\mathcal{D}$, called *quantum differential operators*, are polynomials in the quantities

$$\hbar\partial/\partial t_0, \dots, \hbar\partial/\partial t_r, e^{t_0}, \dots, e^{t_r}, \hbar. \quad (3.6)$$

The $\mathcal{D}$ -module

$$\mathcal{D}(X) = \mathcal{D}/I, \quad I = \{D \in \mathcal{D} \mid DS_a = 0, 1 \leq a \leq r\} \quad (3.7)$$

generated by $S_a = \langle s_a, 1 \rangle$ is called the *Givental quantum $\mathcal{D}$ -module of X* .

Givental and Kim showed that the quantum $\mathcal{D}$ -module of the flag variety G/B is generated by non-constant first integrals of the quantum Toda chain for the Langlands dual $G^\vee$ [12, 20]. Guest and Otofujii [14] generalized some of these results to the case of the affine flag variety $\mathcal{FL}^{(n)}$.

We use this cohomological Givental construction as a motivation and adopt a different strategy in the rest of the paper.

4 Hitchin systems, G -opers and Beilinson-Drinfeld duality

Let X be a smooth projective curve of genus $g \geq 2$ and $\text{Bun}_G(X)$ the moduli space of stable principal G -bundles on X . If $G = \text{GL}_n$ we denote $\text{Bun}_{X,n}$ the moduli space of stable vector bundles of rank n . Hitchin [15] constructed a map

$$h : T^* \text{Bun}_G(X) \rightarrow V \quad (4.1)$$

from the cotangent bundle $T^* \text{Bun}_G(X)$ to a certain vector space V of dimension

$$\dim V = \dim \text{Bun}_G(X) = (g-1) \dim G + \dim Z(G). \quad (4.2)$$

If $G = \text{GL}_n$ then

$$(\text{Bun}_G(X))_P \simeq H^0(X, \text{End } P \otimes K_X) \quad (4.3)$$

where K_X denotes the canonical bundle of X . In other words, $T^* \text{Bun}_{X,n}$ consists of pairs (P, φ) where $P \in \text{Bun}_{X,n}$ and $\varphi : P \rightarrow P \otimes K_X$ a twisted endomorphism. The Hitchin map is defined by

$$h : T^* \text{Bun}_{X,n} \longrightarrow V = \bigoplus_{i=1}^n H^0(X, K_X^i) \quad (4.4)$$

$$(P, \varphi) \longmapsto \text{ch}(\varphi) = (\text{tr } \varphi, \text{tr}(\wedge^2 \varphi), \dots, \text{tr}(\wedge^n \varphi)) \quad (4.5)$$

where $\text{ch}(\varphi)$ is the n -tuple of the coefficients of the characteristic polynomial of φ .

Hitchin has shown that for any basis $\{f_1, \dots, f_r\}$ of V^* the functions $h^* f_i$ and $h^* f_j$ commute with respect to the standard Poisson bracket:

$$\{h^* f_i, h^* f_j\} = 0 \quad (4.6)$$

on $T^* \text{Bun}_G(X)$. Denote by $B_G(X)$ the ring of polynomial functions on V . The Hitchin map induces a morphism

$$\pi_{\text{cl}} : B_G(X) \rightarrow \{\text{functions on } T^* \text{Bun}_G(X)\} \quad (4.7)$$

Suppose now, for the sake of simplicity, that G is semi-simple and simply connected. Beilinson and Drinfeld constructed a *quantization* ($\hbar$ -deformation) of π_{cl} , that is, a graded $\mathbb{C}$ -algebra $A_G(X)$ such that $\text{gr } A_G(X) = B_G(X)$ equipped with a morphism

$$\pi : A_G(X) \longrightarrow H^0(\text{Bun}_G(X), \mathcal{D}^{1/2}) = H^0(\text{Bun}_G(X), \mathcal{D}_{-h^\vee}) \quad (4.8)$$

compatible with π_{cl} . Here $h^\vee$ is the dual Coxeter number of G and $\mathcal{D}^{1/2} = \mathcal{D}_{-h^\vee}$ is the sheaf of twisted differential operators (of critical level $k = -h^\vee$) acting on $K_{\text{Bun}_G(X)}^{1/2} = \eta^{-h^\vee}$, where η^k , $k \in \mathbb{Z}$, denotes a line bundle on $\text{Bun}_G(X)$ obtained via the generalized Borel-Weil-Bott (Kumar-Mathieu) theorem [2, 3, 10].

The elements of $A_G(X)$ are called $G^\vee$ -opers. Denote by $\mathcal{M}(\text{Bun}_G(X))$ (resp. $\mathcal{M}(X \times \text{Bun}_G(X))$) the category of $\mathcal{D}$ -modules on $\text{Bun}_G(X)$ (resp. $X \times \text{Bun}_G(X)$).

THEOREM 4.1 (BEILINSON-DRINFELD) *There exists a family of $\mathcal{D}$ -modules $\{\mathcal{D}_{\mathcal{L}}\}$ on $\text{Bun}_G(X)$ parametrized by $G^\vee$ -opers $\{\mathcal{L}\}$ which are eigensheaves of Hecke operators:*

$$T_\chi : \mathcal{M}(\text{Bun}_G(X)) \longrightarrow \mathcal{M}(X \times \text{Bun}_G(X)), \quad T_\chi \mathcal{D}_{\mathcal{L}} = V_{\mathcal{L}}^\chi \boxtimes \mathcal{D}_{\mathcal{L}} \quad (4.9)$$

for any dominant coweight $\chi \in P_+(G^\vee)$ where $V_{\mathcal{L}}^\chi$ is a $G^\vee$ -local system on X associated to a couple $(\mathcal{L}, \chi)$.

One can give an explicit description of GL_n -opers.

DEFINITION 4.1 A GL_n -oper over X is a local system $(\mathcal{E}, \nabla)$ adapted to a complete flag

$$0 = \mathcal{E}_0 \subset \mathcal{E}_1 \subset \dots \subset \mathcal{E}_n = \mathcal{E} \quad (4.10)$$

of vector bundles over X such that

- (i) $\text{gr}_i(\mathcal{E})$ are invertible,
- (ii) $\nabla(\mathcal{E}_i) \subset \mathcal{E}_{i+1} \otimes K_X$,
- (iii) ∇ induces an isomorphism

$$\text{gr}_i(\mathcal{E}) \xrightarrow{\sim} \text{gr}_{i+1}(\mathcal{E}) \otimes K_X. \quad (4.11)$$

In the local coordinates a GL_n -oper is of the form:

$$\partial_t + \begin{pmatrix} * & * & \cdots & * \\ + & * & \ddots & \vdots \\ & \ddots & \ddots & * \\ 0 & & + & * \end{pmatrix} \quad (4.12)$$

where elements denoted by $*$ are arbitrary and elements denoted by $+$ are non-zero. Locally, one can write a GL_n -oper in the form

$$\partial_t - \begin{pmatrix} q_1 & q_2 & \cdots & q_n \\ 1 & 0 & \cdots & 0 \\ & \ddots & \ddots & \vdots \\ 0 & & 1 & 0 \end{pmatrix}, \quad q_i \in \mathbb{C}[[t]], \quad (4.13)$$

equivalent to a differential operator

$$L = \partial_t^n - q_1 \partial_t^{n-1} - \cdots - q_n. \quad (4.14)$$

Thus, a *local GL_n -oper* is a differential operator of order n whose principal symbol is equal to 1.

5 Affine GL_n -opers and KdV hierarchies

We suppose now that X is a complete (possibly singular) curve and P is a smooth closed point on X .

DEFINITION 5.1 A *local affine GL_n -oper* is a pair $(\widehat{\mathcal{E}}, \widehat{\nabla})$ where $\widehat{\mathcal{E}}$ is a formal torsion-free sheaf of rank n over $X \times \widehat{D}_t = X \times \mathrm{Spec} \mathbb{C}[[t]]$ (an infinitesimal deformation of a torsion-free sheaf $\mathcal{E}$ of rank n over X) equipped with a parabolic structure along of $P \times \widehat{D}_t$:

$$\widehat{\mathcal{E}} = \widehat{\mathcal{E}}_0 \subset \widehat{\mathcal{E}}_1 \subset \cdots \subset \widehat{\mathcal{E}}_n = \widehat{\mathcal{E}}(P) \quad (5.1)$$

Moreover, $\widehat{\nabla}$ is a (micro)connexion on $\widehat{\mathcal{E}}$, that is, a map

$$\widehat{\nabla} : \left(\varinjlim \widehat{\mathcal{E}}_i \right) \longrightarrow \left(\varinjlim \widehat{\mathcal{E}}_i \right) \quad (5.2)$$

satisfying the Leibniz rule

$$\widehat{\nabla}(fs) = f\widehat{\nabla}(s) + \frac{\partial f}{\partial t} \cdot s \quad (5.3)$$

for any function f on $X \times \widehat{D}_t$ and any section s , and such that

- (i) $\mathrm{gr}_i(\widehat{\mathcal{E}})$ are invertible and $\chi(\widehat{\mathcal{E}}_{i+1}) = \chi(\widehat{\mathcal{E}}_i) + 1$,

- (ii) $\widehat{\nabla}(\widehat{\mathcal{E}}_i) \subset \widehat{\mathcal{E}}_{i+1}$ and $\text{ord}_P(\widehat{\nabla}s) = \text{ord}_P(s) + 1$, and
- (iii) $\text{gr}_i(\widehat{\mathcal{E}}) \xrightarrow{\sim} \text{gr}_{i+1}(\widehat{\mathcal{E}})$ is an isomorphism induced by $\widehat{\nabla}$.

PROPOSITION 5.1 (DRINFELD [8]) *There exists a natural bijection between local affine GL_n -opers and Krichever modules, i. e., commutative subrings $R \subset \mathbb{C}[[t]][\partial_t]$, of rank n .*

REMARK 5.1 In positive characteristic, this corresponds to a bijection between elliptic sheaves over a field and Drinfeld modules.

The famous Krichever map [25] associates to any local affine GL_n -oper a point W on the n -reduced Sato Grassmannian $\mathcal{GR}^{(n)}$.

The n th Korteweg-de Vries (KdV) hierarchy describes isospectral deformations of a differential operator $L \in \mathbb{C}[[t]][\partial_t]$ of order n . It may be written in the Lax form:

$$\frac{\partial L}{\partial t_r} = [L_+^{r/n}, L], \quad r \in \mathbb{N}, \quad (5.4)$$

where $L_+^{r/n}$ is the positive part of the microdifferential operator

$$L^{r/n} \in \mathbb{C}[[t]](\partial_t^{-1}). \quad (5.5)$$

Indeed, by a lemma of Schur [26, p. 140], any differential operator $L \in \mathbb{C}[[t]][\partial_t]$ of order n has an (essentially unique) root $L^{1/n} \in \mathbb{C}[[t]](\partial_t^{-1})$.

One can view the n th KdV hierarchy as a dynamical system on the space of local affine GL_n -opers.

6 Semi-infinite q -wedges, quantum τ -functions and quantum $\mathcal{D}$ -modules for KdV theory

Consider the infinite-dimensional Clifford algebra $\mathcal{CL}$ generated by $\psi_n^\pm$, $n \in \mathbb{Z} + 1/2$, satisfying the following relations:

$$[\psi_m^\pm, \psi_n^\pm]_+ = 0, \quad [\psi_m^+, \psi_n^-]_+ = \delta_{m, -n}. \quad (6.1)$$

This Clifford algebra admits a unique irreducible representation (called *spin representation*) in an infinite-dimensional vector space $\mathcal{V}$ (resp. $\mathcal{V}^*$) which is a left (resp. right) module admitting a non-zero *vacuum vector* $|0\rangle$ (resp. $\langle 0|$) such that

$$\psi_i^\pm |0\rangle = 0 \quad (\text{resp. } \langle 0| \psi_{-i}^\pm = 0) \quad \text{for } i > 0. \quad (6.2)$$

Denote by $\mathfrak{gl}_\infty$ the Lie algebra of infinite matrices indexed by $\mathbb{Z}+1/2$ with almost all entries equal to zero. It has a canonical basis consisting of matrices E_{ij} , $i, j \in \mathbb{Z}+1/2$, with 1 as the (i, j) -entry and 0 elsewhere. The map $E_{ij} \mapsto \psi_{-i}^+ \psi_j^-$ defines a bijection between $\mathfrak{gl}_\infty$ and quadratic elements $\sum a_{ij} \psi_{-i}^+ \psi_j^-$ of $\mathcal{CL}$. Later $\mathfrak{gl}_\infty$ will also be identified with normally ordered expressions of the type $\sum a_{ij} : \psi_{-i}^+ \psi_j^- :$.

Using the exponential map we obtain a representation R of the group GL_∞ on $\mathcal{V}$ and $\mathcal{V}^*$. For a positive integer m denote

$$\langle \pm m | = \langle 0 | \psi_{1/2}^\pm \cdots \psi_{m-1/2}^\pm \in \mathcal{V}^* \text{ and } | \pm m \rangle = \psi_{-m+1/2}^\pm \cdots \psi_{-1/2}^\pm | 0 \rangle \in \mathcal{V}.$$

This defines the *charge decomposition* $\mathcal{V} = \bigoplus \mathcal{V}^{(m)}$ (cf. below).

This spin representation has the following semi-infinite wedge realization. Let $\mathbb{C}^\infty$ be an infinite-dimensional complex vector space with a fixed basis $\{v_i\}_{i \in \mathbb{Z}+1/2}$. Then the *semi-infinite wedge* (or *fermionic Fock space*) $\mathcal{V} = \wedge^{\infty/2} \mathbb{C}^\infty$ is generated by semi-infinite monomials of the form

$$\begin{aligned} v_{i_1} \wedge v_{i_2} \wedge v_{i_3} \wedge \cdots \quad \text{where } i_1 > i_2 > i_3 > \cdots \\ \text{and } i_{k+1} = i_k - 1 \text{ for } k \gg 0. \end{aligned} \tag{6.3}$$

The generators of the infinite-dimensional Clifford algebra are represented as the wedging and contracting operators defined by:

$$\begin{aligned} \psi_j^+(v_{i_1} \wedge v_{i_2} \wedge \cdots) &= v_{-j} \wedge v_{i_1} \wedge v_{i_2} \wedge \cdots \\ &= (-1)^s v_{i_1} \wedge \cdots \wedge v_{i_s} \wedge v_{-j} \wedge v_{i_{s+1}} \cdots \quad \text{if } i_s > -j > i_{s+1} \\ \psi_j^-(v_{i_1} \wedge v_{i_2} \wedge \cdots) &= (-1)^{s+1} v_{i_1} \wedge \cdots \wedge v_{i_{s-1}} \wedge v_{i_{s+1}} \cdots \quad \text{if } j = i_s \end{aligned}$$

and zero in all other cases. By definition, the *charge* of $v_{i_1} \wedge v_{i_2} \wedge \cdots$ is equal to m if $i_k + k = m + 1/2$ for $k \gg 0$. This defines the charge decomposition as above. In particular,

$$|m\rangle = v_{m-1/2} \wedge v_{m-3/2} \wedge v_{m-5/2} \cdots \in \mathcal{V}^{(m)}. \tag{6.4}$$

The space $\mathcal{V}^{(m)}$ is an irreducible highest weight representation of $\mathfrak{gl}_\infty \subset \mathcal{CL}$ with the highest weight vector $|m\rangle$.

The GL_∞ -orbit of the vacuum vector $|0\rangle$ is closely related to the infinite-dimensional Sato Grassmannian. Indeed, let us identify $\mathbb{C}^\infty$ with $\mathbb{C}((t))$ via $v_i \leftrightarrow t^{i+1/2}$. Then the map

$$\varphi : \mathrm{GL}_\infty |0\rangle \longrightarrow \mathcal{GR}, \quad \tau = \bigwedge_{\{i=-1/2, -3/2, \dots\}} u_i \longmapsto \sum_{\{i=-1/2, -3/2, \dots\}} \mathbb{C} u_i$$

is well-defined since $u_{-i} = v_{-i}$ for i big enough. This map is surjective and we can also write

$$\mathcal{GR} = \mathrm{GL}_\infty |0\rangle / \mathbb{C}^*. \tag{6.5}$$

By definition, semi-infinite wedges lying in the $\mathbf{GL}_\infty$ -orbit of $|0\rangle$ are called τ -functions in the fermionic picture.

Consider the *affine Hecke* (or *Iwahori-Hecke*) algebra $\widehat{\mathcal{H}}_N(q)$ with the generators $T_i^{\pm 1}$, $X_i^{\pm 1}$, $1 \leq i \leq N-1$, and relations:

$$T_i T_i^{-1} = T_i^{-1} T_i = X_i X_i^{-1} = X_i^{-1} X_i = 1; \quad (6.6a)$$

$$T_i T_j = T_j T_i \text{ if } |i-j| > 1, \quad X_i X_j = X_j X_i \quad \forall i, j; \quad (6.6b)$$

$$T_i T_{i+1} T_i = T_{i+1} T_i T_{i+1}, \quad (T_i + 1)(T_i - q) = 0; \quad (6.6c)$$

$$X_j T_i = T_i X_j \text{ if } j \neq i, i+1, \quad T_i X_i T_i = q X_{i+1}. \quad (6.6d)$$

Its infinite counterpart $\widehat{\mathcal{H}}_\infty(q)$ is generated by a countable number of generators $T_i^{\pm 1}$ and $X_i^{\pm 1}$, $i \in \mathbb{N}$, satisfying the same relations.

Let V be a complex vector space of dimension n with a basis $\{e_1, \dots, e_n\}$ and let $V(\mathfrak{z}) = V \otimes_{\mathbb{C}} \mathbb{C}[\mathfrak{z}, \mathfrak{z}^{-1}]$. We identify $V(\mathfrak{z})$ with $\mathbb{C}^\infty$ via the correspondence $\mathfrak{z}^j e_i \leftrightarrow v_{i-nj-1/2}$. There are well-known right actions of $\widehat{\mathcal{H}}_N(q^2)$ on $V(\mathfrak{z})^{\otimes N}$ and of $\widehat{\mathcal{H}}_\infty(q^2)$ on the so-called *thermodynamic limit* $V(\mathfrak{z})^{\otimes \infty/2}$ [19, 28]. The latter is well-defined since any T_i acts only in a pair of adjacent factors. The q -antisymmetrization procedure (loc. cit.) gives the q -wedge spaces

$$\bigwedge_q^N V(\mathfrak{z}) = V(\mathfrak{z})^{\otimes N} \bigg/ \sum_{i=1}^{N-1} \text{Ker}(T_i + 1) \quad (6.7a)$$

and

$$\bigwedge_q^{\infty/2} V(\mathfrak{z}) = V(\mathfrak{z})^{\otimes \infty/2} \bigg/ \sum_{i=1}^{\infty} \text{Ker}(T_i + 1). \quad (6.7b)$$

In the same manner, the q -deformed Fock space $\mathcal{V}_q^{(m)}$ of charge m is the quotient of the space $U^{(m)}$ of pure tensors $u_{m_1} \otimes u_{m_2} \otimes \dots$ where $m_k = m - k + 1/2$ for $k \gg 1$ by $\sum_{i=1}^{\infty} (T_i + 1)$.

DEFINITION 6.1 The elements of the q -antisymmetrized $\mathbf{GL}_\infty$ -orbit of the vacuum vector $|0\rangle$ will be called *quantum τ -functions*.

Using fermionic normal ordering

$$:\psi_m \psi_n^*: = \begin{cases} \psi_m \psi_n^* & \text{if } m < 0 \text{ or if } n > 0, \\ -\psi_n^* \psi_m & \text{if } m > 0 \text{ or if } n < 0 \end{cases} \quad (6.8)$$

one can define operators

$$H_n = \sum_{i \in \mathbb{Z} + 1/2} :\psi_{-i}^+ \psi_{i+n}^-: \quad (6.9)$$

satisfying the commutation relations:

$$[H_m, H_n] = m\delta_{m,-n}. \quad (6.10)$$

It is easy to see that for any given element u of the Fock space, the expression $H_n|u\rangle$ is a finite sum. Consider also the generating function

$$H(t) = \sum_{n=1}^{\infty} t_n H_n \quad (6.11)$$

called the *Hamiltonian* where $t = (t_1, t_2, \dots)$. In the bosonic picture, the τ -function τ_g , associated to $g = \sum a_{ij} : \psi_{-i}^+ \psi_j^- : \in \mathfrak{gl}_{\infty}$, is the correlation function

$$\tau_g(t) = \langle 0 | e^{H(t)} g | 0 \rangle = \langle 0 | e^{H(t)} e^{\sum a_{ij} : \psi_{-i}^+ \psi_j^- :} | 0 \rangle \quad (6.12)$$

(*cf.* [6, 7, 16]). The wedging and contracting operators ψ_i^+ and ψ_j^- correspond to the intertwiners:

$$\Gamma_i^+(\mathfrak{z}) : V(\mathfrak{z}) \otimes \mathcal{V}^{(m)} \longrightarrow \mathcal{V}^{(m+1)} \text{ and } \Gamma_j^-(\mathfrak{z}) : \mathcal{V}^{(m)} \longrightarrow \mathcal{V}^{(m-1)} \otimes V(\mathfrak{z}). \quad (6.13)$$

Now let W be a point of $\mathcal{GR}$ associated to a local affine $\mathbf{GL}_n$ -oper $(\widehat{\mathcal{E}}, \widehat{\nabla})$ via the Krichever map (*cf.* sect. 5). Consider the associated τ -function

$$\tau_W(t) = \langle 0 | e^{H(t)} g_W | 0 \rangle = \langle 0 | e^{H(t)} e^{\sum a_{ij} : \psi_{-i}^+ \psi_j^- :} | 0 \rangle \quad (6.14)$$

The wedging and contracting operators ψ_i^+ and ψ_j^- correspond to the intertwiners of level-1 $\widehat{\mathfrak{sl}}_n$ -modules

$$\Gamma_i^+(\mathfrak{z}) : V(\mathfrak{z}) \otimes \mathcal{V}_{\Lambda_m} \longrightarrow \mathcal{V}_{\Lambda_{m+1}} \text{ and } \Gamma_j^-(\mathfrak{z}) : \mathcal{V}_{\Lambda_m} \longrightarrow \mathcal{V}_{\Lambda_{m-1}} \otimes V(\mathfrak{z}) \quad (6.15)$$

where $\mathcal{V}_{\Lambda_m}$ are the $\widehat{\mathfrak{sl}}_n$ -modules with fundamental weights Λ_m .

In general, consider a *singular vector* of a highest weight $\widehat{\mathfrak{sl}}_n$ -module V_{Λ} . It is an eigenvector of the Cartan subalgebra of $\widehat{\mathfrak{sl}}_n$ annihilated by $U_+(\widehat{\mathfrak{sl}}_n)$ different from the highest weight vector v_{Λ} . It may be written in the form Sv_{Λ} where S is a uniquely defined element of $U_-(\widehat{\mathfrak{sl}}_n)$. Then

$$\langle 0 | (Sv_{\Lambda})g | 0 \rangle = D \langle 0 | v_{\Lambda}g | 0 \rangle = 0. \quad (6.16)$$

for a certain differential operator D [9, sect. 5.3].

In our case, any τ -function τ_W defines a $\mathcal{D}$ -module $\mathcal{D}(\tau_W)$ on $\mathcal{GR}$. These $\mathcal{D}$ -modules are closely related to the so-called deformed Knizhnik-Zamolodchikov equations for form factors (*cf.* [27, sect. 6]).

REMARK 6.1 All τ -functions satisfy bilinear Hirota equations (of KP hierarchy) related to the time evolution $e^{H(t)}$. These equations characterize the Sato Grassmannian itself.

Some remarks can be made, however, in our infinite-dimensional setting. The ind-varieties $\mathcal{GR}$ and $\mathcal{GR}^{(n)}$ have two equivalent ind-structures given by pole orders and by Schubert varieties [21]. Denote by $\mathcal{GR} = \varinjlim \mathcal{GR}_i$ one of these structures and let $\mathcal{D}(\mathcal{GR}) = \varinjlim \mathcal{D}(\mathcal{GR}_i)$ be the sheaf of differential operators on $\mathcal{GR}$. Then one considers right $\mathcal{D} = \mathcal{D}(\mathcal{GR})$ -modules in the abelian category $\mathcal{M}(\mathcal{GR}, \mathcal{O})$ of $\mathcal{O}^!$ -modules (*cf.* [2, sect. 7.11] for a detailed exposition).

Quantum correlations functions constructed from intertwiners of $U_q(\widehat{\mathfrak{sl}}_n)$ -modules should satisfy quantum differential equations in the sense of Frenkel-Reshetikhin [11]. We use the notation $\mathcal{D}_q(\tau_W)$ for associated *quantum* $\mathcal{D}$ -modules on $\mathcal{GR}^{(n)}$ parametrized by affine local GL_n -opers.

7 Affine Miura opers and Miura transformation

We use [4, 7.2 and 8.3.5] as a reference for affine Miura opers. Let G be a reductive group, X a smooth complex curve and $P \in |X|$. Denote $\mathrm{LG}_X^- \subset \mathrm{LG}$ the subgroup consisting of loops which extends to $X \setminus P$.

Let B be a Borel subgroup of G and B^- a transverse Borel subgroup (such that $[B^-] \in G/B$ lies in the open B -orbit). Choose a point Q on X distinct from P and consider the subgroup $\mathrm{LG}_X^{--} \subset \mathrm{LG}_X^-$ consisting of “negative” loops whose values at Q lie in B^- .

DEFINITION 7.1 A *local affine Miura G -oper* is a quadruple $(\mathcal{U}, \nabla, \mathcal{U}^{--}, \mathcal{U}^+)$ where $\mathcal{U}$ is an LG -torsor on $\widehat{D}_t$ with a connection ∇ , a flat reduction $\mathcal{U}^{--}$ to LG_X^{--} and a reduction $\mathcal{U}^+$ to LG^+ in “tautological relative position” with respect to ∇ (*loc. cit.*).

REMARK 7.1 One can define local affine G -opers in the same vein replacing LG_X^{--} by LG_X^- .

Local Miura GL_n -oper is a connection of the type

$$\partial_t - \begin{pmatrix} \chi_1 & 0 & 0 & \cdots & 0 \\ 1 & \chi_2 & 0 & \cdots & 0 \\ 0 & 1 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & 1 & \chi_n \end{pmatrix}, \quad \chi_i \in \mathbb{C}[[t]], \quad (7.1)$$

on a vector bundle of rank n over $\widehat{D}_t$. The *Miura transformation* is the transformation from this connection to the unique gauge equivalent connection of the form (4.13). Geometrically, it may be described as follows [10, 3.8]. Let $(\mathcal{E}, \nabla)$ be a local GL_n -oper with another full flag of subbundles

$$\mathcal{E}'_1 \subset \mathcal{E}'_2 \cdots \subset \mathcal{E}'_n \quad (7.2)$$

preserved by ∇ . Such a flag defines a connection ∇' on $\text{gr}(\mathcal{E}) = \oplus_{i=1}^n \text{gr}_i(\mathcal{E})$. Then the Miura transformation associates $(\mathcal{E}, \nabla)$ to $(\text{gr}(\mathcal{E}'), \nabla')$.

Now let $(\widehat{\mathcal{E}}, \widehat{\nabla})$ be a local affine GL_n -oper and

$$\widehat{\mathcal{E}} = \widehat{\mathcal{E}}'_0 \subset \widehat{\mathcal{E}}'_1 \subset \widehat{\mathcal{E}}'_2 \subset \cdots \subset \widehat{\mathcal{E}}'_n = \widehat{\mathcal{E}}'(Q) \quad (7.3)$$

another parabolic structure preserved by $\widehat{\nabla}$. As above, the *affine Miura transformation* associates $(\widehat{\mathcal{E}}, \widehat{\nabla})$ to $(\text{gr}(\widehat{\mathcal{E}}'), \widehat{\nabla}')$.

PROPOSITION 7.1 *Let $(\text{gr}(\widehat{\mathcal{E}}'), \widehat{\nabla}')$ be a local affine Miura GL_n -oper and $(\widehat{\mathcal{E}}, \widehat{\nabla})$ the associated local affine GL_n -oper. Let W be a point of the n -reduced Sato Grassmannian $\mathcal{GR}^{(n)}$ which is the image of $(\widehat{\mathcal{E}}, \widehat{\nabla})$ under the Krichever map. Then*

$$(\text{gr}(\widehat{\mathcal{E}}'), \widehat{\nabla}') \mapsto (W, \widehat{\mathcal{E}}'_1/\widehat{\mathcal{E}}' \subset \widehat{\mathcal{E}}'_2/\widehat{\mathcal{E}}' \subset \cdots \subset \widehat{\mathcal{E}}'_n/\widehat{\mathcal{E}}') \quad (7.4)$$

defines a point of the affine flag variety $\mathcal{FL}^{(n)}$.

Proof. Recall that there is a natural fibration $\mathcal{FL}^{(n)} \rightarrow \mathcal{GR}^{(n)}$ whose fiber is the usual flag variety GL_n/B . Thus, the pair

$$(W, \widehat{\mathcal{E}}'_\bullet/\widehat{\mathcal{E}}') = (W, \widehat{\mathcal{E}}'_1/\widehat{\mathcal{E}}' \subset \widehat{\mathcal{E}}'_2/\widehat{\mathcal{E}}' \subset \cdots \subset \widehat{\mathcal{E}}'_n/\widehat{\mathcal{E}}') \quad (7.5)$$

defines, indeed, a point of $\mathcal{FL}^{(n)}$ under this fibration.

8 $\mathcal{D}$ -modules for affine Toda theories

Consider the generating functions

$$\psi^+(\mathfrak{z}) = \sum_{i \in \mathbb{Z}} \psi_i^+ \mathfrak{z}^i \text{ and } \psi^-(\mathfrak{z}) = \sum_{i \in \mathbb{Z}} \psi_i^- \mathfrak{z}^{-i} \quad (8.1)$$

of free fermions $\psi_i^\pm$ indexed by $i \in \mathbb{Z}$ (cf. sect. 6). Introduce “positive” and “negative” times $t = (t_1, t_2, \dots)$ and $t' = (t'_1, t'_2, \dots)$ resp., and denote $\xi(t, \mathfrak{z}) = \sum_{n \geq 0} t_n \mathfrak{z}^n$. The *two-dimensional Toda theory* is related to the following time evolution $g(t, t')$:

$$\psi^+(\mathfrak{z}) \mapsto \mathfrak{z}^n e^{\xi(t, \mathfrak{z}) + \xi(t', \mathfrak{z}^{-1})} \psi(\mathfrak{z}) \quad (8.2)$$

$$\psi^-(\mathfrak{z}) \mapsto \mathfrak{z}^{-n} e^{-\xi(t, \mathfrak{z}) - \xi(t', \mathfrak{z}^{-1})} \psi(\mathfrak{z}) \quad (8.3)$$

(with essential singularities at $\mathfrak{z} = 0, \infty$) of $g = \exp(\sum_i a_i \psi^+(p_i) \psi^-(q_i))$ [16, §9]. The τ -functions are defined as before by

$$\tau_g(t, t') = \langle 0 | g(t, t') | 0 \rangle \quad (8.4)$$

(loc. cit.). The usual reduction to $A_n^{(1)}$ [16, §8,9] defines τ -functions of affine Toda chains.

Let $(\text{gr } (\widehat{\mathcal{E}}'), \widehat{\nabla}')$ be a local affine Miura GL_n -oper and denote $(W, \widehat{\mathcal{E}}'_\bullet / \widehat{\mathcal{E}}')$ the associated point of $\mathcal{FL}^{(n)}$. For any τ -function

$$\tau_W(t, t') = \langle 0 | g_W(t, t') | 0 \rangle \quad (8.5)$$

corresponding to $(W, \widehat{\mathcal{E}}'_\bullet / \widehat{\mathcal{E}}')$, we construct $\mathcal{D}$ -modules $\mathcal{D}(\tau_W)$ and $\mathcal{D}_q(\tau_W)$ as in sect. 6.

9 Proof of the main theorem

The result follows easily from the constructions above.

Let $(\text{gr } (\widehat{\mathcal{E}}'), \widehat{\nabla}')$ be a local affine Miura GL_n -oper and $(W, \widehat{\mathcal{E}}'_\bullet / \widehat{\mathcal{E}}')$ the associated point of $\mathcal{FL}^{(n)}$. Under the Miura transformation, we get a local affine GL_n -oper $(\mathcal{E}, \nabla)$. It is easy to see that the Miura transform of $(W, \widehat{\mathcal{E}}'_\bullet / \widehat{\mathcal{E}}')$ coincides with the point $W \in \mathcal{GR}^{(n)}$ obtained from $(\mathcal{E}, \nabla)$ via the Krichever map.

Moreover, the τ -function $\tau_W(t, t')$ of the affine Toda $\widehat{\mathfrak{sl}}_n$ -chain becomes the τ -function $\tau_W(t)$ of the n th KdV hierarchy. Clearly, this transformation preserves null vectors. Thus, the Givental type $\mathcal{D}$ -modules $\mathcal{D}(\tau_W)$ and $\mathcal{D}_q(\tau_W)$ on $\mathcal{FL}^{(n)}$ are transformed to Beilinson-Drinfeld type $\mathcal{D}$ -modules on $\mathcal{GR}^{(n)}$.

10 Alternative construction of $\mathcal{D}$ -modules

In this section we briefly indicate how one can construct $\mathcal{D}$ -modules on $\mathcal{GR}^{(n)}$ and $\mathcal{FL}^{(n)}$ using a local affine version of the Beilinson-Drinfeld approach.

In the case of the n th KdV hierarchy, an analogue of the Hitchin fibration is the fibration $S \rightarrow \mathcal{AO}^{(n)}$ whose fibers are (Jacobians) of spectral curves. Here $\mathcal{AO}^{(n)}$ denotes the moduli space of local affine GL_n -opers (so-called *abelianized Grassmannian*). Recall (cf. [1, 22]) that any local affine GL_n -oper $m \in \mathcal{M}$, i. e., any Krichever module, defines a spectral curve X_m . The fibration above is closely related to $T^*\mathcal{GR}^{(n)}$ and one can construct Beilinson-Drinfeld type $\mathcal{D}$ -modules on $\mathcal{GR}^{(n)}$ indexed by local affine GL_n -opers.

In the case of the affine Toda $\widehat{\mathfrak{sl}}_n$ -chain, an analogue of the Hitchin fibration is the fibration $S \rightarrow \mathcal{MO}^{(n)}$ whose fibers are (Jacobians) of spectral curves of this chain [24, sect. 1]. Here $\mathcal{MO}^{(n)}$ denotes the moduli space of local affine Miura GL_n -opers. Using the Beilinson-Drinfeld procedure one can construct $\mathcal{D}$ -modules on $\mathcal{FL}^{(n)}$ indexed by such opers.

Acknowledgements. I am grateful to Mark Spivakovsky for useful remarks.

References

- [1] M. R. Adams and M. J. Bergvelt, *The Krichever map, vector bundles over algebraic curves and Heisenberg algebras*, Comm. Math. Phys. **154** (1993), 1–60.
- [2] A. A. Beilinson and V. G. Drinfeld, *Quantization of Hitchin’s integrable system and Hecke eigensheaves*, draft version on the D. Ben-Zvi homepage at <http://zaphod.uchicago.edu/>.
- [3] ———, *Quantization of Hitchin’s fibration and Langlands program*, Algebraic and geometric methods in mathematical physics (Kaciveli, 1993) (A. Boutet de Monvel and V. A. Marchenko, eds.), Math. Phys. Stud., no. 19, Kluwer Acad. Publ., Dordrecht, 1996, pp. 3–7.
- [4] D. Ben-Zvi and E. Frenkel, *Spectral curves, opers and integrable systems*, arXiv:math.AG/9902068, 64 p.
- [5] D. A. Cox and Sh. Katz, *Mirror symmetry and algebraic geometry*, Mathematical Surveys and Monographs, vol. 68, American Mathematical Society, Providence, RI, 1999.
- [6] E. Date, M. Jimbo, M. Kashiwara, and T. Miwa, *Transformation groups for soliton equations*, Nonlinear integrable systems – classical theory and quantum theory (Kyoto, 1981) (M. Jimbo and T. Miwa, eds.), World Scientific, Singapore, 1983, pp. 39–120.
- [7] E. Date, M. Jimbo, and T. Miwa, *Solitons. Differential equations, symmetries and infinite-dimensional algebras*, Cambridge Tracts in Mathematics, vol. 135, Cambridge University Press, Cambridge, 2000.
- [8] V. G. Drinfeld, *Commutative subrings of certain noncommutative rings (Russian)*, Funkcional. Anal. i Priložen **11** (1977), no. 1, 11–14, English translation: Funct. Anal. Appl. **11** (1977), no. 1, 9–12.
- [9] B. Feigin and F. Malikov, *Modular functor and representation theory of $\widehat{\mathfrak{sl}}_2$ at a rational level*, Operads: Proceedings of Renaissance Conferences (Hartford, CT/Luminy, 1995), Contemp. Math., no. 202, Amer. Math. Soc., Providence, RI, 1997, (arXiv:q-alg/9511011), pp. 357–405.
- [10] E. Frenkel, *Affine algebras, Langlands duality and Bethe ansatz*, XIth International Congress of Mathematical Physics (Paris, 1994), Internat. Press, Cambridge, 1995, (arXiv:q-alg/9506003), pp. 606–642.
- [11] I. B. Frenkel and N. Yu. Reshetikhin, *Quantum affine algebras and holonomic difference equations*, Comm. Math. Phys. **146** (1992), no. 1, 1–60.
- [12] A. Givental, *Stationary phase integrals, quantum Toda lattices, flag manifolds and the mirror conjecture*, Topics in singularity theory, Amer. Math. Soc. Transl. Ser. 2, no. 180, Amer. Math. Soc., Providence, RI, 1997, (arXiv:alg-geom/9612001), pp. 103–115.

- [13] A. Givental and B. Kim, *Quantum cohomology of flag manifolds and Toda lattices*, Comm. Math. Phys. **168** (1995), no. 3, 609–641, arXiv:hep-th/9312096.
- [14] M. Guest and T. Otofujii, *Quantum cohomology and periodic Toda lattice*, Comm. Math. Phys. **217** (2001), no. 3, 475–487, arXiv:math.QA/9812127.
- [15] N. Hitchin, *Stable bundles and integrable systems*, Duke Math J. **54** (1987), no. 1, 91–114.
- [16] M. Jimbo and T. Miwa, *Solitons and infinite-dimensional Lie algebras*, Publ. Res. Inst. Math. Sci. **19** (1983), no. 3, 943–1001.
- [17] V. G. Kac and A. K. Raina, *Bombay lectures on highest weight representations of infinite-dimensional Lie algebras*, Advanced Series in Mathematical Physics, vol. 2, World Scientific, Teaneck, NJ, 1987.
- [18] V. G. Kac and J. W. van de Leur, *The n -component KP hierarchy and representation theory*, Important Developments in Soliton Theory, Springer Ser. Nonlinear Dynam., Springer, Berlin, 1993, pp. 302–343.
- [19] M. Kashiwara, T. Miwa, and E. Stern, *Decomposition of q -deformed Fock spaces*, Selecta Math. (N.S.) **1** (1995), no. 4, 787–805, arXiv:q-alg/9508006.
- [20] B. Kim, *Quantum cohomology of flag manifolds G/B and quantum Toda lattices*, Ann. of Math. (2) **149** (1999), no. 1, 129–148.
- [21] S. Kumar, *Infinite Grassmannians and moduli space of G -bundles*, Vector bundles on curves – new directions (Cetraro, 1995), Lecture Notes in Math., no. 1649, Springer, Berlin, 1997, pp. 1–49.
- [22] Yi. Li and M. Mulase, *Hitchin systems and KP equations*, Internat. J. Math. **7** (1996), no. 2, 227–244.
- [23] Yu. I. Manin, *Frobenius manifolds, quantum cohomology, and moduli spaces*, AMS Colloquium Publications, vol. 47, American Mathematical Society, Providence, RI, 1999.
- [24] A. McDaniel, *Representations of $\mathfrak{sl}(n, \mathbb{C})$ and the Toda lattice*, Duke Math. J. **56** (1988), no. 1, 47–99.
- [25] M. Mulase, *Category of vector bundles on algebraic curves and infinite-dimensional Grassmannians*, Internat. J. Math. **1** (1990), no. 3, 293–342.
- [26] D. Mumford, *An algebro-geometric construction of commuting operators and of solutions of the Toda lattice equation, Korteweg-de Vries equation and related non-linear equations*, Proceedings of the International Symposium on Algebraic Geometry (Kyoto Univ., Kyoto, 1977), Kinokuniya Book Store, Tokyo, 1978, pp. 115–153.

- [27] F. A. Smirnov, *Form factors, deformed Knizhnik-Zamolodchikov equations and finite-gap integration*, Comm. Math. Phys. **155** (1993), no. 3, 459–487, arXiv:hep-th/9210052.
- [28] E. Stern, *Semi-infinite wedges and vertex operators*, Internat. Math. Res. Notices (1995), no. 4, 201–219, arXiv:q-alg/9504013.
- [29] A. Strominger, Sh.-T. Yau, and E. Zaslow, *Mirror symmetry is T-duality*, Nuclear Phys. B **479** (1996), no. 1-2, 243–259, arXiv:hep-th/9606040.
- [30] C. Vafa and E. Witten, *A strong coupling test of S-duality*, Nuclear Phys. B **431** (1994), no. 1-2, 3–77, arXiv:hep-th/9408074.

Igor Potemine
 Laboratoire Emile Picard
 Université Paul Sabatier
 118, route de Narbonne
 31062 Toulouse Cédex

e-mail : potemine@picard.ups-tlse.fr